

Searches for Techniparticles at DØ

Lorenzo Feligioni

Boston University

for the DØ collaboration

- Technicolor Phenomenology
- Signatures:
 - $W \pi_T \rightarrow e \nu b \bar{b}/\bar{c}$
 - $\omega_T/\rho_T \rightarrow e^+e^-$
- Cross Section Limits



Technicolor Physics

- Mechanism of electroweak symmetry breaking
 - Standard Model and Supersymmetry:
 - Higgs field $\Rightarrow$ Higgs boson
 - Weakly coupled, light fundamental scalar
 - Technicolor:
 - New strong dynamics: $SU(N_{TC})$ gauge theory analog to QCD
 - $N_{TC}^2 - 1$ new gauge bosons: technigluons
 - In analogy with QCD breaking of the chiral symmetry produces Goldstone bosons (technipions):
 - 3 technipions are eaten to become W_L and Z_L
 - others could be observed at collider experiments



Low Scale Technicolor Models

- Large numbers of technifermions are the natural choice for several Technicolor Models
 - **Walking Technicolor**
 - Evade large flavor changing neutral current
 - **Topcolor-assisted Technicolor**
 - Many technifermions are needed to generate hard masses for quarks and leptons
 - **Technicolor Straw Man Model (TCSM2):** K. Lane, S. Mrenna hep-ph/02110299
 - Set the scale for calculating lowest-lying bound state of lightest technifermion doublets
 - color singlet vectors (200 – 400 GeV)
 - produced in pp collisions

» Decays:

$\omega_T \rightarrow \gamma \pi_T$	$\rho_T \rightarrow \pi_T \pi_T$	$\rho_T \rightarrow \pi_T \pi_T$
$\rightarrow \gamma Z$	$\rightarrow W \pi_T, Z \pi_T$	$\rightarrow W \pi_T, Z \pi_T$
$\rightarrow 3 \pi_T$	$\rightarrow WW$	$\rightarrow WZ$
$\rightarrow f f, g g$	$\rightarrow f f, g g$	

- color-singlet scalars
 - lightest technihadrons $\pi_T^0 \pi_T^{+/-}$
 - » Decays:

$$\pi_T \rightarrow ff, gg \quad (\pi_T^0 \rightarrow bb, \pi_T^{+/-} \rightarrow bc \text{ dominate})$$



Previous Searches

- TCSM2 Parameters:
 - N_D : number of technifermion doublets
 - $Q_D = Q_U - 1$: technifermion charge
 - $\sin\chi$: mixing angle
 - M_V : mass parameter (it controls technifermions coupling and decay mode)

- Previous searches

CDF RunI



- $W\pi_T$ and $\omega_T \rightarrow \gamma\pi_T$
- $M_V = 100$

DØ RunI


- $\rho_T/\omega_T \rightarrow ee$
- $M_T = M_V = 100$ to 400
- $M(\rho_T/\omega_T) - M(\pi_T) = 60, 100$ GeV
 - $M_V=100$ GeV $\Rightarrow W\pi_T$ channel open

LEP



- $\rho_T \rightarrow WW, \rho_T \rightarrow \pi_T W$ (DELPHI)
- $M(\pi_T) = 105$ GeV $M(\rho_T)=200$ GeV is excluded for some TCSM parameters



$W\pi_T$ Events Selection



- **One reconstructed Primary Vertex**
- **One isolated electron**
 - veto on the presence of another electrons suppress Z contamination
- **Missing $E_T > 20$ GeV**
 - eliminates multi-jets (QCD)
- **Two calorimeter jets**
 - Veto on a third jet, suppresses $t\bar{t}$ background
 - At least one jet has to be associated with a **Secondary Vertex (b-tagging)**
- **$M_T(W) > 30$ GeV**
- **E_T^W ($p_T(e) + \text{missing } E_T$) > 65 GeV**







$W\pi_T$ Searches

$M(\pi_T) = 105 \text{ GeV}$ $M(\rho_T) = 200 \text{ GeV}$

Cross sections ($W \rightarrow e\nu$), $M_V = 200$:

- $W\pi_T^{+/-} \sim 3.7 \text{ pb}$
- $W\pi_T^0 \sim 2.9 \text{ pb}$

$$\int \mathcal{L} dt = 238 \text{ pb}^{-1}$$



Data

74

Sources of Background

Physics Background

tt	15
Single top	4
W+ Heavy Flavors	33
WZ	1
Z→ee	2
total	55

Instrumental Background

QCD	7
W + light quarks	11

Tot Backd

73 ± 19

Expected Signal

9.1 ± 1.3



DPF 30 August 04

Lorenzo Feligioni





6



$W\pi_T$ Optimization

$$\int \mathcal{L} dt = 238 \text{ pb}^{-1}$$

- H_T^e (electron $p_T + \sum \text{jet } p_T$)
- $p_T(jj)$ (p_T of the dijet system)
- $\Delta\phi(jj)$
- $M(jj)$ (invariant mass of the dijet system)
- $M(Wjj)$ (invariant mass of the W + dijet system)




$W\pi_T$ Cross Section Limit

- $\Delta\phi(jj) > 2.2$
- $p_T(jj) > 75$ GeV
- $H_T^e < 200$ GeV
- Mass Window

$$\int \mathcal{L} dt = 238 \text{ pb}^{-1}$$

data background signal

Baseline + $\Delta\phi$	28	28.3 ± 7.1	7.5 ± 1.1
+ $p_T(jj)$	22	24.7 ± 6.2	7.4 ± 1.1
+ H_T^e	17	18.3 ± 4.6	7.2 ± 1.1
+ mass window	4	6.6 ± 1.6	6.2 ± 0.9



Cross section 95% C.L. upper limit 6.4 pb



DPF 30 August 04

Lorenzo Feligioni


8



$$\omega_T/\rho_T \rightarrow e^+e^-$$

$$\int \mathcal{L} dt = 200 \text{ pb}^{-1}$$

- Events with 2 high p_T electrons are selected
- Background
 - Drell-Yan production
 - QCD
- Search for $\rho_T/\omega_T \rightarrow e^+e^-$ as a bump/excess at high dielectron mass
- Intrinsic widths of ρ_T, ω_T are about 0.5 GeV
 - Thus resonance width dominated by detector resolution



$\omega_T/\rho_T \rightarrow e^+e^-$ Limits

$$\int \mathcal{L} dt = 200 \text{ pb}^{-1}$$


❖ 95% C.L.

➤ $m(\rho_T, \omega_T) - m(\pi_T) = 60 \text{ GeV}$

- $m(\rho_T, \omega_T) > 367 \text{ GeV}$
for $M_V = 500 \text{ GeV}$
- $m(\rho_T, \omega_T) > 340 \text{ GeV}$
for $M_V = 100 \text{ GeV}$

➤ $m(\rho_T, \omega_T) - m(\pi_T) = 100 \text{ GeV}$

- $m(\rho_T, \omega_T) > 355 \text{ GeV}$
for $M_V = 500 \text{ GeV}$
- $m(\rho_T, \omega_T) > 240 \text{ GeV}$
for $M_V = 100 \text{ GeV}$



Summary

- DØ has begun to search for Technicolor particles in the $W+2$ jets channel
 - New b-tagging capability respect to RunI
 - No evidence were found for the π_T , ρ_T mass combination considered
- $\rho_T/\omega_T \rightarrow ee$ analysis
 - Most restrictive constraints on dilepton technicolor decays to date
- Outlook:
 - Almost twice more luminosity available for these analysis
 - Add μ channels soon

