



Inclusive W and Z cross sections at the Tevatron

Victoria Martin

Northwestern University

Of behalf of the CDF and DØ collaborations



Meeting of The Division of Particles and Fields
of the APS, 2004

Contents

- What's so interesting about inclusive $W^\pm$, Z^0 cross sections?
- Fermilab Tevatron, DØ and CDF detectors
- Reconstruction, efficiencies, backgrounds
- Systematic uncertainties
- Estimating PDF uncertainties
- Cross Section Results
- R and $\Gamma(W)$
- Future Plans and Conclusions

Fermilab Tevatron and Luminosity

- $\sqrt{s}=1960$ TeV $p \bar{p}$ collider
- Upgraded in 2000
- Two collision points: CDF and DØ
- Luminosity uncertainty is 6%. Limited by:
 - Knowledge of inelastic cross section
 - Acceptance of luminosity counters



- Current luminosity: 680 pb^{-1}
- Luminosity on tape: $\sim 500 \text{ pb}^{-1}$
- Measurements presented here: $\sim 60 \text{ pb}^{-1}$ to $\sim 200 \text{ pb}^{-1}$

Why W and Z Physics at RunII?

- Large numbers of W bosons
- High mass Z/γ^* events
- Sensitive to u & d quarks

New Forward Detectors

Di-boson cross sections

Tri-boson couplings

Beyond SM:

Di-bosons couplings

Z' resonance

anomalous q couplings

W, Z cross sections

Z Forward-Backward Asymmetry

Indirect $\Gamma(W)$

W asymmetry

$\sin^2(\theta_W)$,
quark couplings

Lepton Universality

W mass

Direct $\Gamma(W)$


Higgs Mass
Constraint

Cross Section Calculation


$$\sigma \cdot \text{BR} = \frac{N_{\text{cand}} - N_{\text{backg}}}{L \cdot A \cdot \varepsilon}$$

- N_{cand} : *Number of candidate events*
- N_{backg} : *Estimated number of background events*
- L : *Integrated luminosity*
- $A \cdot \varepsilon$: *Acceptance X efficiency*
- Look for W and Z decays to leptonic final states
 - need to reconstruct e, μ , τ , ν
 - requires good calorimeter, muon and tracking coverage

CDF Run II Detector



DØ Detector



Backgrounds

- 3 main background sources:

- QCD processes


- jet looks like a lepton
 - signal events tend to have isolated ℓ
 - ℓ is further from other detector activity
 - look at isolated and non-isolated events

- Other EW processes

- calculate from MC



- Cosmics (in μ channel)

- require small cda between μ and beam spot (CDF/DØ)
 - reconstruct cosmic tracks (CDF)
 - residual background very small (<1%)



Acceptance and Efficiency

- Geometric acceptance calculated from Pythia or Herwig MC
- Use NLO CTEQ5 PDFs
- All other efficiencies calculated from data
 - Many efficiencies measured from $Z \rightarrow \ell + \ell^-$ events
 - 1st ℓ is identified, test the 2nd ℓ




$W \rightarrow \ell \nu$ Inclusive Cross Section

- $W \rightarrow e \nu$ reconstruction:

- missing- $E_T > 25 \text{ GeV}$
- CDF: 1 central electron ($|\eta| < 1.0$) or 1 forward electron ($1.0 < |\eta| < 3.2$)
- with matching track
- DØ: 1 central electron ($|\eta| < 1.05$)
- no track requirement

- $W \rightarrow \mu \nu$ reconstruction:

- missing- $E_T > 20 \text{ GeV}$
- 1 muon ($|\eta| < 0.6$) with track




$$M_T = \sqrt{(E_T(\ell) + E_T(\nu))^2 - (p_x(\ell) + p_x(\nu))^2 - (p_y(\ell) + p_y(\nu))^2}$$

		Events	Background	$A \cdot \epsilon$
CDF central $W \rightarrow e \nu$	72 pb^{-1}	37584	$(4.4 \pm 0.8)\%$	$(18.0 \pm 0.4)\%$
CDF forward $W \rightarrow e \nu$	64 pb^{-1}	10461	$(8.7 \pm 0.6)\%$	$(5.2 \pm 0.2)\%$
CDF $W \rightarrow \mu \nu$	194 pb^{-1}	57109	$(7.3 \pm 0.5)\%$	$(9.6 \pm 0.2)\%$
DØ $W \rightarrow e \nu$	177 pb^{-1}	175572	$(33.6 \pm 0.4)\%$	$(22.6 \pm 0.5)\%$

$Z \rightarrow e^+e^-$ results



- CDF reconstruction:
 - 2 central electrons ($|\eta| < 1.0$), both with matching tracks
 - 1 central electron ($|\eta| < 1.0$) with matching track, plus 1 forward electron ($1.0 < |\eta| < 3.2$)
 - $66 < m(ee)/\text{GeVc}^{-2} < 116$
- DØ reconstruction:
 - 2 central electrons ($|\eta| < 1.05$)
 - at least 1 matching track
 - $70 < m(ee)/\text{GeVc}^{-2} < 110$
- Correct σ for DY



	Events	$A \cdot \epsilon$
CDF 72pb ⁻¹	4242	(22.7±0.5)%
DØ 177pb ⁻¹	4712	(9.6±0.4)%

$Z \rightarrow \mu^+ \mu^-$ results




- CDF reconstruction:
 - 1 central muon ($|\eta| < 0.6$) with matching track
 - 1 additional central track ($|\eta| < 1.0$)
 - $66 < m(\mu\mu)/\text{GeV}c^{-2} < 116$
- DØ reconstruction:
 - 2 muons, both matched to tracks
 - $m(\mu\mu) > 40 \text{ GeV}/c^2$
- <2% background
- Correct σ for DY



	Events	$A \cdot \epsilon$
CDF 194pb ⁻¹	3568	(7.3±0.2)%
DØ 148pb ⁻¹	14352	22.1% to 31.6% (t-dep)

DØ: reconstructing τ

- Use Neural Net to separate τ from background
 - NN detects $\tau \rightarrow \text{hadrons } \nu_\tau$, $\tau \rightarrow e \nu_e \nu_\tau$
 - input: EM energy and track p_T distributions
 - trained on MC, cut $\text{NN} > 0.8$





• $Z \rightarrow \tau^+ \tau^-$ signal

- 1 NN τ and 1 $\tau \rightarrow \mu \nu \nu$
- 1946 events found in 207 pb^{-1}
- $(55 \pm 2)\%$ background from misidentified τ 's
- $A \cdot \varepsilon = (1.65 \pm 0.09)\%$


CDF: τ results

- Look for hadronic tau decays
 - Narrow isolated jet
 - Low track multiplicity
 - mass of (tracks, π^0) $< m(\tau)$
- $Z \rightarrow \tau^+ \tau^-$ signal
 - 1 hadronic τ and 1 $\tau \rightarrow e \nu \nu$ decay



	events	$A \cdot \epsilon$	backg
$Z \rightarrow \tau^+ \tau^-$ 72pb ⁻¹	50	~0.5%	(21 $\pm$ 6)%
W $\rightarrow \tau \nu$ 72pb ⁻¹	2345	(0.91 $\pm$ 0.05)%	(26 $\pm$ 3)%



Estimating PDF Uncertainties



- PDF uncertainty on acceptance is largest systematic
- CTEQ suggested method
- NNLO CTEQ6L provides:
 - default PDFs
 - 20 pairs of error PDFs, corresponding to varying each parameter in 90% CL interval
- Use numerical integration to calculate change in acceptance for pair
- MRST error sets also checked

Cross Section Results

- Black: stat + syst error
- Aqua: luminosity error
- Yellow: NNLO prediction



Inclusive Cross Sections Results

CDF (central) $\sigma \cdot \text{BR}(W \rightarrow e \nu) = 2780 \pm 14(\text{stat}) \pm 60(\text{sys}) \pm 167(\text{lum}) \text{ pb}$

CDF (forward) $\sigma \cdot \text{BR}(W \rightarrow e \nu) = 2874 \pm 34(\text{stat}) \pm 167(\text{sys}) \pm 172(\text{lum}) \text{ pb}$

DØ $\sigma \cdot \text{BR}(W \rightarrow e \nu) = 2865 \pm 8(\text{stat}) \pm 63(\text{sys}) \pm 40(\text{pdf}) \pm 186(\text{lum}) \text{ pb}$

CDF $\sigma \cdot \text{BR}(W \rightarrow \mu \nu) = 2768 \pm 12(\text{stat}) \pm {}^{+65}_{-55}(\text{sys}) \pm 166(\text{lum}) \text{ pb}$

CDF $\sigma \cdot \text{BR}(W \rightarrow \tau \nu) = 2620 \pm 70(\text{stat}) \pm 210(\text{sys}) \pm 160(\text{lum}) \text{ pb}$

NNLO prediction

Stirling, Van Neerven

$\sigma \cdot \text{BR}(p\bar{p} \rightarrow W \rightarrow \ell \nu) = 2687 \pm 54 \text{ pb}$

CDF $\sigma \cdot \text{BR}(Z \rightarrow e^+ e^-) = 255.8 \pm 3.9(\text{stat}) \pm 5.5(\text{sys}) \pm 15.3(\text{lum}) \text{ pb}$

DØ $\sigma \cdot \text{BR}(Z \rightarrow e^+ e^-) = 255.8 \pm 3.9(\text{stat}) \pm 8.5(\text{sys}) \pm 5.1(\text{pdf}) \pm 15.3(\text{lum}) \text{ pb}$

CDF $\sigma \cdot \text{BR}(Z \rightarrow \mu^+ \mu^-) = 253.1 \pm 4.2(\text{stat}) \pm {}^{+8.3}_{-6.4}(\text{sys}) \pm 14.9(\text{lum}) \text{ pb}$

DØ $\sigma \cdot \text{BR}(Z \rightarrow \mu^+ \mu^-) = 291.3 \pm 3.0(\text{stat}) \pm 6.9(\text{sys}) \pm 18.9(\text{lum}) \text{ pb}$

CDF $\sigma \cdot \text{BR}(Z \rightarrow \tau^+ \tau^-) = 242 \pm 49(\text{stat}) \pm 31(\text{sys}) \pm 16(\text{lum}) \text{ pb}$

DØ $\sigma \cdot \text{BR}(Z \rightarrow \tau^+ \tau^-) = 256 \pm 16(\text{stat}) \pm 17(\text{sys}) \pm 16(\text{lum}) \text{ pb}$

NNLO prediction

Stirling, Van Neerven

$\sigma \cdot \text{BR}(p\bar{p} \rightarrow Z \rightarrow \ell \ell) = 253.1 \pm 5.0 \text{ pb}$

Cross Section Ratio, R

$$R = \frac{\sigma(p\bar{p} \rightarrow W \rightarrow \ell \nu)}{\sigma(p\bar{p} \rightarrow Z \rightarrow \ell \ell)}$$

- Ratio of W and Z cross sections
 - Luminosity, and many systematics cancel

CDF
(72 pb⁻¹ central e, μ combined) $R = 10.92 \pm 0.15(stat) \pm 0.14(sys)$

DØ (e) $R = 10.82 \pm 0.16(stat) \pm 0.25(sys) \pm 0.13(pdf)$

NNLO prediction $R = 10.69 \pm 0.08$

BR(W → ℓν) and Γ(W)

CDF:

$$R = \frac{\sigma(p\bar{p} \rightarrow W \rightarrow \ell \nu)}{\sigma(p\bar{p} \rightarrow Z \rightarrow \ell \ell)} = 10.92 \pm 0.15(\text{stat}) \pm 0.14(\text{sys})$$

$$R = \frac{\sigma(p\bar{p} \rightarrow W)}{\sigma(p\bar{p} \rightarrow Z)} \cdot \frac{\Gamma(Z)}{\Gamma(Z \rightarrow \ell \ell)} \cdot \frac{\Gamma(W \rightarrow \ell \nu)}{\Gamma(W)}$$

3.3677 ± 0.024
NNLO (PDG)


From LEP:
(3.366 ± 0.002)%

$$\text{BR}(W \rightarrow \ell \nu) = 0.1089 \pm 0.0022$$

Using NNLO calculation

Γ(W → ℓν) = 226.4 ± 0.3 MeV (PDG):

$$\Gamma(W) = 2079 \pm 41 \text{ MeV}$$



Current World Average: 2124 ± 41 MeV



Conclusions & prospects for more data

- We've measured inclusive W and Z production at the Tevatron in all 6 leptonic channels.
- More improvements will be made on R and $\Gamma(W)$
- **Combine CDF and DØ results!**
- Cross section measurements in e and μ channels are already systematics limited
 - Some systematic errors will reduce with statistics
 - Main uncertainty comes from PDF
- **Think smarter! Select events in such a way to cancel errors**
- **Use W and Z cross sections for normalization?**
 - Measurements have ~2% uncertainty
 - Luminosity error is 6%

Backup Slides

PDF Uncertainties II

- Acceptance as a function of boson rapidity



$$A_W \propto 4 \int_0^{y_{\max}} x_a x_b u(x_a) d(x_b) A_W(y) dy$$

$$x_{a,b} = \frac{M_W}{\sqrt{s}} \exp(\pm y)$$

$$A_Z \propto \int_0^{y_{\max}} x_a x_b (C_u u(x_a) u(x_b) + C_d d(x_a) d(x_b)) A_Z(y) dy$$

Luminosity Measurement at CDF

- Luminosity (L) from Cherenkov Luminosity Counters:
- Discriminating primary (from ppbar)-secondary particles
- Time resolution

→ Well adapted for high-L collider

$$\mu_{\alpha} \cdot f_{BC} = \sigma_{pp} \cdot \epsilon_{\alpha}^{clc} \cdot L$$

α : selection criteria for pp coll.

μ_a : #int/Bunch Crossing (BC)

f_{BC} : BC Frequency

σ_{pp} : $p\bar{p}$ inelastic cross section

ϵ_{α}^{clc} : Efficiency of CLC

Increasing Lum.

"0" hits

#hits

#part

$$\mu_{clc} = -\ln\left(\frac{N_{zeroBC}}{N_{totalBC}}\right)$$

$$\mu_{clc} = \frac{\langle N_{Hits/BC} \rangle}{\langle N_{Hits/p\bar{p}} \rangle}$$

$$\mu_{clc} = \frac{\sum A_i}{\langle A_{p\bar{p}} \rangle}$$

Dominant uncertainties from:

e_a = MC simulation.


$\sigma_{pp} = 60.4 \pm 1.4$ mb (measured by CDF)

μ_{clc} : Different definitions

Systematic uncertainty ~5%

Neutrino reconstruction

- CDF and DØ are hermetic transverse to the beam
- Colliding partons have no transverse momentum
- Measure neutrinos by the amount of energy missing transverse to the beam
- For $W \rightarrow \ell \nu$ candidates calculate transverse mass (M_T) from x and y components



$$M_T = \sqrt{(E_T(\ell) + E_T(\nu))^2 - (p_x(\ell) + p_x(\nu))^2 - (p_y(\ell) + p_y(\nu))^2}$$

Systematic Uncertainties

- One example: $D\bar{D} W \rightarrow e\nu$, $Z \rightarrow e^+e^-$ cross section

Source	Relative Uncertainty (%) on		
	σ_W	σ_Z	R
stat			
Number of Events	0.29	1.47	1.50
sys			
Stat. Error on Efficiencies	0.72	0.77	0.58
Background Subtraction	0.35	0.82	0.47
Fake Track Match Prob.	0.45	n/a	0.45
Monte Carlo Parameters	1.13	0.84	1.41
Trigger Eff	0.59	0.06	0.53
Electron ID	1.43	2.86	1.43
Track Match	0.52	0.35	0.17
Total sys	2.19	3.21	2.27
pdf			
PDF	1.41	1.94	1.21
lumi			
Luminosity	6.5	6.5	n/a

Cross sections already
systematic limited

Largest error from
PDFs, ID cuts