

Top Quark Mass Measurement **in the lepton+jets channel using** ***Template Method* and** ***Dynamical Likelihood Method* at CDF**

Kohei Yorita

Waseda University, Japan
For the CDF collaboration

Aug.30th 2004


APS/DPF @ Riverside CA



Introduction

- ▶ From RunI results : New World Average (2004)
 $M_t = 178.0 \pm 4.3 \text{ GeV } (\pm 2.7 \pm 3.3) \text{ hep-ex/0404010}$
- ▶ Top quark is heavy ($\sim 180 \text{ GeV}$)
 Yukawa coupling ~ 1 .

$$y_t = \frac{\sqrt{2} m_t}{v} \approx 1$$
- * The mass is near the Electro-Weak Symmetry Breaking Scale.
- ▶ Correlated with Higgs mass, together with W boson mass.




$$dM_W \propto (M_{\text{top}}^2, \ln(M_H))$$

- ▶ Standard Model Higgs Mass:
 Most probable: $114 \pm_{45}^{69} \text{ GeV}$
 Upper limit(95% CL): 260 GeV
- ▶ In this talk,
 - (1) Dynamical Likelihood Method (DLM)
 lepton+jets ³ 1b tag
 - (2) Template Method
 lepton+jets 0tag, ³ 1tag, $\longrightarrow$ combined

RunII Goal : $\pm 2 \sim 3 \text{ GeV}$!

Event Selection




► “lepton+jets” decay mode

Final State : lepton(e,m), n, 4jets(2bjets)

► SVX b jet tagging

Identify by finding a
vertex of displaced tracks

Excellent silicon detector!



Cuts	DLM	Template(btag)	Template(0tag)
Lepton/MET	One central electron or muon($E_t(P_t) > 20\text{GeV}$), $\text{Met} > 20\text{GeV}$		
nJets	$= 4\text{jets}(E_t > 15)$	$\geq 3.5\text{jets}(E_t > 15, > 8)$	$\geq 4\text{jets}(E_t > 21)$
b tag	$\geq 1\text{ btag}$	$\geq 1\text{ btag}$	NO
kinematical fit	No	YES ($c^2 < 9$)	YES ($c^2 < 9$)
Observed	22 ev ($L = 162\text{pb}^{-1}$)	28 ev ($L = 162\text{pb}^{-1}$)	39 ev ($L = 194\text{pb}^{-1}$)
S/B	~ 4.2	~ 3.0	~ 0.7



Dynamical Likelihood Method



- Originally proposed in 1988 by K.Kondo.(J.Phys. Soc. 57, 4126)

► **For i -th event, likelihood is defined as,**

$$L^i(M_{top}) = \sum_{I_t} \sum_{I_s} \int \frac{2p^4}{Flux} F(z_a, z_b, p_T) |M|^2 d(s_w - (\ell + n)^2) w(I_t, \mathbf{x} | \mathbf{y}; M_{top}) d\mathbf{x} ds_w$$

F : Parton distribution function for (z_a, z_b) and Pt of tt system

M : Matrix element of tt process, $|M|^2 = |M_{prod}|^2 \Pi(s_r) |M_{dec}|^2$

w : Transfer function, $\mathbf{x}$; partons $\longleftrightarrow$ $\mathbf{y}$; observables

$$s_w = (l+v)^2$$

► **Two Summations and one integration :**

I_t : Jets to partons combination (6: 1-tag, 2: 2-tag), I_s : Two v_z solutions,
In practice, integration of $\mathbf{x}$, s_w made by Monte Carlo Method.

► **Extract top mass by maximum likelihood method,**


$$\Lambda(M_{top}) = -2 \ln \left(\prod_{event} L^i(M_{top}) \right) \longrightarrow \overline{\mathbf{M}}_{top} = \mathbf{M}_{top} \text{ min. } \Lambda(M_{top})$$



Background Correction : DLM



- ▶ In the method, first, all events are assumed to be signal. Need to correct background effect.
- ▶ Backgrounds pull likelihood peak down. (Have Lower Mass)
- ▶ Pseudo experiments(22ev) by varying background fraction to look at effects on signal tt events.





- ▶ Use this map to extract the top mass.

source	W+4j
Mistag	1.2 ± 0.37
Wbb	0.7 ± 0.29
Wcc	0.3 ± 0.12
Wc	0.2 ± 0.12
Single top	0.17 ± 0.03
WW	0.08 ± 0.05
nonW	1.6 ± 0.38
Bkg tot.	4.2 ± 0.71
N obs.	22
tt (6.7pb)	20.9



Pull Distribution : DLM

After applying mapping functions




- No bias was found. The statistical uncertainties are also scaled by the slope of mapping function properly.


Results : DLM

- Observed events : Total 22 events; electrons 12, Muons 10

22 events joint likelihood



Correct background-pulling
4.2 events expected.





Statistical & Systematic Uncertainty



Expected statistical Error


Black arrows :

Data : + 4.5, - 5.0 GeV

Monte Carlo :

Mean : + 5.4, - 5.0 GeV

MPV : + 4.5, - 4.1 GeV



DLM : $M_{\text{top}} = 177.8 \pm 4.5$ (stat.) ± 6.2 (syst.) GeV/c²

Systematic uncertainty

Sources	delta M_{top} (GeV/c ²)
Jet Energy Scale	5.3
Transfer function	2.0
PDF	2.0
Generator	0.6
MC Modeling	0.6
ISR	0.5
FSR	0.5
Bkg fraction ($\pm 5\%$)	0.5
Bkg Modeling	0.5
Spin correlation	0.4
NLO effect	0.4
Total	6.2 GeV

(NLO: re-weighted qq and gg fractions)




Template Method Overview

(1) For each event, invariant mass of top is reconstructed by minimizing χ^2 defined as,

$$\chi^2 = \sum_{i=\ell, \text{4 jets}} \frac{(p_T^{i, \text{fit}} - p_T^{i, \text{meas}})^2}{\sigma_i^2} + \sum_{j=x, y} \frac{(p_j^{UE, \text{fit}} - p_j^{UE, \text{meas}})^2}{\sigma_j^2} + \frac{(M_{jj} - M_W)^2}{\Gamma_W^2} + \frac{(M_{\ell\nu} - M_W)^2}{\Gamma_W^2} + \frac{(M_{bjj} - M_t)^2}{\Gamma_t^2} + \frac{(M_{b\ell\nu} - M_t)^2}{\Gamma_t^2}$$

- Jets resolution term
- Unclustered Energy term
- W mass and Top mass term

Top mass is free parameter




(2) Additional selection cut on resulting χ^2

(3) Build templates of M_t with smallest χ^2 from MC for

- Signal with different top mass
- Each background source

(4) Parameterization : Build signal p.d.f. as a function of generated mass.


(5) Fit to the data using unbinned likelihood.






Template Method with ≥ 1 b tag

CDF Run II Preliminary (162 pb⁻¹)



Constrained fit

CDF Run II Preliminary (162 pb⁻¹)



Unconstrained fit

Category	Background (events)		M_{top} (GeV/c ²)
	input	output	
$W + \geq 3.5$ jets	7.0 ± 0.8	7.1 ± 0.8	$174.9^{+6.7}_{-7.2}$
	no constraint	$16.7^{+8.2}_{-7.3}$	$177.7^{+9.3}_{-8.4}$

Template (btag) : $M_{\text{top}} = 174.9 \pm 7.1^{+7.1}_{-7.7}$ (stat.) ± 6.5 (syst.) GeV/c²


Stat. error from the fit is scaled up by 1.065 to provide 68% coverage of the true mass value

Jet Energy Scale : 6.3 GeV




Template Method with 0tag

CDF II Preliminary (193.5 pb⁻¹)



CDF II Preliminary (193.5 pb⁻¹)



Input (signal)	Output	M_{top}
15.5 ± 3.2	16 ± 3	$179.1 \pm \begin{matrix} 10.5 \\ 9.5 \end{matrix}$
No constraint	26 ± 13	$177.5 \pm \begin{matrix} 9.1 \\ 7.7 \end{matrix}$

Template (0tag) : $M_{\text{top}} = 179.1 \pm \begin{matrix} 10.5 \\ 9.5 \end{matrix}$ (stat.) ± 8.4 (syst.) GeV/c²




Jet Energy Scale : 8.3 GeV

- ▶ Divided into 4 sub-samples :
 - 2 tag (4 combinations)
 - 1 tag (12 combinations)
 - 3.5 jets, 4jets different S/B
 - 0 tag, (24 combinations)

- ▶ Mass templates built for each sub-samples.
- ▶ Likelihood procedure uses information of all sub-samples simultaneously to determine best top mass.

$$L = L_{2tag} \times L_{1tag,4j} \times L_{1tag,3.5j} \times L_{0tag}$$

- ▶ Uncertainty is scaled up by 1.058 for 68% error coverage.




Results : Template Method


Combined likelihood curve

$$L = L_{2tag} \times L_{1tag,4j} \times L_{1tag,3.5j} \times L_{0tag}$$

Likelihood vs top mass



Expected Statistical Uncert.



CDF Run II Preliminary

Source of Systematics	$\Delta M_{\text{top}} (\text{GeV}/c^2)$
Jet Energy Scale	6.8
ISR	1.2
FSR	1.2
PDFs	0.4
Generators	0.3
Background Shape	1.0
Other MC modeling	0.8
b tagging	0.1
MC statistics	0.3
Total	7.1

$$\text{Template : } M_{\text{top}} = 176.7 \pm \begin{matrix} 6.0 \\ 5.4 \end{matrix} \text{ (stat.)} \pm 7.1 \text{ (syst.) GeV}/c^2$$



Summary



DLM : (Current CDF official)

$$M_{\text{top}} = 177.8 \pm 4.5 \text{ (stat.)} \pm 6.2 \text{ (syst.) GeV/c}^2$$

Combined Template : (less assumption, RunI check)

$$M_{\text{top}} = 176.7 \pm 6.0 \text{ (stat.)} \pm 7.1 \text{ (syst.) GeV/c}^2$$

(1) Now our measurements are dominated by systematics.


- Reduce Jet Energy Scale Uncert. by a better understanding of cal. simulation.

(2) Add more data (expect : >40' RunI in RunII)
(now : 1.5' RunI)

(3) Measurement of hadronic W mass
- constrain light jet energy scale


(4) Will try to combine diff. methods,
diff. channels to get one value!

Again, RunII goal is an error of 2~3 GeV





Back up slides

Some Plots (Data vs MC)




- The data are well understood by MC!
- Agreement is quite good!

(H_t : Et scalar sum of lepton , jets, Met)



- **Event-by-event Maximum Likelihood Mass with wide range of [125-225] GeV.**

Note : First(last) bin includes under(over) flow.



- **Event likelihood**

For i-th event,

$$L_{ev}^i = \int L^i(M) dM$$

➡ A event has one likelihood value

